# Adding Polynomials

To add polynomials, this is VERY similar to collecting like terms, you:

- 1. Drop the brackets we are allowed to do this when there is only a PLUS sign between the brackets \* this does not work with a subtract sign.
- 2. Identify the like terms
- 3. Rearrange (optional) \*remember the sign (+/-) stays with the term
- 4. Add the coefficients \*remember the sign (+/-) stays with the term
- 5. Keep the variable the same


Example 1:	Example 2:
$(2x^2 + 3x + 5) + (x^2 + 2x + 3)$	$(4y^2 - 2y - 5) + (-y^2 + 3y + 3)$
$= 2x^2 + 3x + 5 + x^2 + 2x + 3$	$=4y^2 - 2y - 5 + -y^2 + 3y + 3$
$= 2x^{2} + x^{2} + 3x + 2x + 5 + 3$ = $3x^{2} + 5x + 8$	$= 4y^2 - 2y - 5 - y^2 + 3y + 3$
	$=3y^2+y-2$

#### **Practice: Adding Polynomials**

a. 
$$(a+1) + (a+1)$$
  
b.  $(2a+3) + (-6a+2)$ 

c. 
$$(4n^2 + 3n + 1) + (n^2 + n + 2)$$
  
d.  $(-p^2 - 2p + 4) + (3p^2 - 2p - 1)$ 

e. Find the 'algebraic expression' for the perimeter of the following triangle.



## Subtracting Polynomials

<b>Finding the opposite:</b> What is the opposite of +5?	? What is the opposite of -7?	
What is the opposite of x?	What is the opposite of -3	λ <sub>5</sub>
Write the opposites of the f	ollowing expressions:	
a5x + 4	b. 6x – y	c. x + y

#### To subtract polynomials, you:

CANNOT drop the brackets! If you drop the brackets, only the first term of the second bracket will be subtracted  $\rightarrow$  the entire bracket following the minus sign needs to be subtracted.

 We need a + between the brackets in order to remove the brackets. We can change the – to a +, if we also change everything in the following bracket to 'the opposite'. This is known as ADDING THE OPPOSITE (the additive inverse).

Then it is the same as adding polynomials!

- 2. Drop the brackets we are allowed to do this when there is only a PLUS sign between the brackets \* this does not work with a subtract sign.
- 3. Identify the like terms
- 4. Rearrange (optional) \*remember the sign (+/-) stays with the term
- 5. Add the coefficients \*remember the sign (+/-) stays with the term
- 6. Keep the variable the same

Example 1	Example 2
$(2x^2 + 3x + 5) - (x^2 + 2x + 3)$	$(4y^2 - 2y - 5) - (-y^2 + 3y + 3)$
$= 2x^2 + 3x + 5 + (-x^2 - 2x - 3)$	$=4y^2 - 2y - 5 + (+y^2 - 3y - 3)$
$= 2x^2 + 3x + 5 - x^2 - 2x - 3$	$=4y^2 - 2y - 5 + y^2 - 3y - 3$
$= 2x^2 - x^2 + 3x - 2x + 5 - 3$	=5y <sup>2</sup> -5 y - 8
$= x^2 + x + 2$	

#### **Practice: Subtracting Polynomials**


a. 
$$(a+5) - (2a+1)$$
  
b.  $(2a+3) - (4a+2)$ 

c. 
$$(n^2 + 3n + 1) - (n^2 + n + 2)$$
  
d.  $(-p^2 - 2p + 4) - (3p^2 - 2p - 1)$ 

e. 
$$(3m+3) - (4m-2)$$
  
f.  $(4g^2 - g + 7) - (-2g - 4)$ 

**q.** 
$$(3m^2 - 2m) - (5m + 9) + 4m$$
  
**h.**  $-(m + 7) - (3m + 9)$ 

i. Find an algebraic expression for the length of AB in the following diagram.



### **ANSWERS** a) -a +4, b) -2a + 1, c) 2n - 1, d) -4p<sup>2</sup> + 5, e) -m +5, f) 4g<sup>2</sup> +g +11, g) 3m<sup>2</sup> - 3m - 9, h) -4m -16, i) x + 4