SNC2D CHEMISTRY

CHEMICAL REACTIONS

Conservation of Mass (P.176-177)

Activity: Measuring Mass (Part 1)

INSTRUCTIONS (CLOSED SYSTEM)

- A. Measure \sim 30 mL of vinegar (acetic acid) into a small Erlenmeyer flask.
- B. Using a scoopula, place a small amount of baking soda (sodium hydrogen carbonate powder) into a small balloon.
- C. Carefully place the balloon over the mouth of the Erlenmeyer flask, ensuring the vinegar and baking soda do **not** mix.
- D. Using a balance, measure and record the total mass of the assembly.
- E. Tip the balloon up and empty it's contents into the Erlenmeyer flask. Ensure the balloon does fly off. Record your observations.
- $\mathsf{F}.$ Once the reaction has stopped, measure and record the mass of the assembly again.
- G. Clean up your work area but remain where you are.

February 23, 2013

2DCHEM - Conservation of Mass

Activity: Measuring Mass (Part 1)			
OUESTIONS			
 What evidence of a chemical change did you notice? 			
2. Compare the total mass of the Erlenmeyer flask/balloon assembly before and after the reaction. What do you notice?			
 bubbles of gas are produced, change is difficult to reverse total mass of assembly before and after should be relatively the same 			
February 23, 2013 2DCHEM - Conservation of Mass 2			

Activity: Measuring Mass (Part 2)

INSTRUCTIONS (OPEN SYSTEM)

- H. Fill a small test tube ~ half full with of vinegar (acetic acid).
- I. Using a scoopula, place a small amount of baking soda (sodium hydrogen carbonate powder) into a small beaker.
- Carefully place the test tube into the beaker, ensuring the vinegar and baking soda do **not** mix.
- K. Using a balance, measure and record the total mass of the assembly.
- L. Empty the contents of the test tube into the beaker. Ensure that the reaction does not spill over. Record your observations.
- M. Once the reaction has stopped, place the test tube back into the beaker and then measure and record the mass of the assembly again.N. Clean up your work area and put your equipment away.

February 23, 2013

2DCHEM - Conservation of Mass

3

4

Activity: Measuring Mass (Part 2)

QUESTIONS

- 3. Compare the total mass of the beaker/test tube assembly before and after the reaction. What do you notice?
- 4. Compare the results of the two experiments. What do you notice? Explain any difference.
- 3. total mass before should be greater than the total mass after
- 4. in the closed system mass was conserved in the open system mass was lost the gas lost must have mass

February 23, 2013

2DCHEM - Conservation of Mass

Chemical Reactions

The most important advancement in understanding what happens during chemical reactions was made about two centuries ago by the French chemist Antoine Lavoisier, with the assistance of his wife and research colleague Marie-Anne Paulze. The couple made meticulous measurements of the masses of reactants and products in many kinds of chemical reactions.



February 23, 2013

2DCHEM - Conservation of Mass

.

Chemical Reactions

What they discovered was that the total mass of reactants and the total mass of products in a given reaction are always the same. Another way of saying this is that the mass is conserved (i.e. the mass does not change during a chemical reaction). No known exceptions to this have ever been observed. For this reason, this experimental result has come be known as the **law of conservation** of mass.



6

7

February 23, 2013

2DCHEM - Conservation of Mass

Law of Conservation of Mass

NOTE!

The law of conservation of mass is very important in understanding what happens to the atoms in chemical reactions. It implies that no atoms are destroyed and no new atoms are produced during a chemical reaction. Instead, the atoms in the reactants of a chemical reaction are simply rearranged to form the products. Chemical bonds between atoms are broken and new ones are formed, and the atoms simply reconnect in new ways.


February 23, 2013

2DCHEM - Conservation of Mass





The rearrangement of atoms that occurs during a chemical reaction can be illustrated using models or diagrams. Consider a vehicle that runs on electricity produced in a fuel cell. The electricity comes from a reaction between hydrogen gas and oxygen gas to form liquid water. The equations for this reaction are: word equation: hydrogen + oxygen → water chemical equation: + $O_{2(g)}$ + $H_2O_{(l)}$ H_{2(g)} H2 H₂ 02 H₂0 HoC 2DCHEM - Conservation of Mass February 23, 2013 8

















Law of Conservation of Mass - DYK?

The law of conservation of mass has implications far beyond the laboratory. Think about engines that use fuels as a source of energy. When cars burn gasoline, energy is released during the chemical reaction of combustion. However, since mass is conserved, all of the mass of the fuel is still present in some form after combustion occurs.



February 23, 2013

2DCHEM - Conservation of Mass

Law of Conservation of Mass – DYK?


NOTE!

Every kilogram of fuel mixed with oxygen in an internal combustion engine produces 3 to 4 kg of water vapour and carbon dioxide gas, as well as pollutant gases such as nitrogen oxides and unburned hydrocarbons.



February 23, 2013

2DCHEM - Conservation of Mass



Check Your Learning				
(b) H	/hy does bread rise? ow do you think the mass of the bread would compare to the riginal total mass of the ingredients in the recipe?	2		
	ne yeast reacts with the sugar – produces carbon dioxide nould be slightly lighter – gases have escaped			
February 23, 2	2013 2DCHEM - Conservation of Mass 1:	7		

Check Your Learning	
TEXTBOOK	
P.177 Q.3-5	
February 23, 2013 2DCHEM - Conservation of Mass 18	